

THAWING CANIPLAS

Plasvacc recommends that the following instructions be strictly followed when thawing a bag of Caniplas Canine Plasma.

It must be remembered that any heat above body temperature has the ability to activate the Fibrinogen in the plasma, converting it into Fibrin – in other words, the whole bag turns into a CLOT!

The remaining plasma is still safe to transfuse, however the filter in the giving set will block, and need to be replaced repeatedly.

PREPARATION INSTRUCTIONS:

- I. Gently cut the tamperproof seal on the plasma box
- 2. Working over a bench, preferably covered with some sort of cushion (a doubled towel), tip the bag of Caniplas out of the box, into your hand
- 3. Fill a sink or bucket with water NO WARMER than 40° Centigrade (35°C is ideal)
- 4. Ensure that the hot water is evenly mixed with the cold water
- 5. A thermometer MUST be used
- 6. Please do not use heating elements
- 7. Place the bag into the water have enough water in the sink so that the bag FLOATS
- 8. The thermometer must be in the water bath for the duration of the thawing process

- 9. Further hot water will need to be added during the thawing process to ensure that the plasma is approximately at body temperature for transfusion. The Bag of Caniplas should be removed from the water bath while the extra hot water is added. The plasma cannot come into contact with the hot water (greater than 40° Centigrade) as it is running from the tap
- 10. If you are heating the plasma after thawing it in a fridge, the same rule applies. The water is to be no warmer than 40°C, and a thermometer must be used
- II. If thawing is performed by lay staff, please review their procedures from time to time
- 12. Please thaw straight from the freezer to the warm water bath.

REMEMBER... If the bag of Caniplas is thawed and warmed over a period of less than 20 minutes, too much heat has been used.

Some fine flakes of Fibrin may be visible as the bag thaws, especially while it still contains an "ice block" of frozen plasma. These flakes should disappear as the plasma warms, but it is not unusual if some flakes are still visible at the time of transfusion - their presence is completely normal, and the reason why a blood giving set with a 200 micron filter must be used.

PLASVACC

Rigby 6066 Cunningham Hwy Kalbar Queensland 4309 Phone | 1800 | 7|5|2|7|6|2|

(07) 5463 7770 Fax plasmail@plasvacc.com Email Web www.plasvacc.com

Technical support and inquiries to Plasvacc

Phone 1800 752 762 Email plasmail@plasvacc.com To order

Please contact your local veterinary wholesale supplier

